

Management of *Tuta absoluta*

1



IPM INNOVATION LAB
Feed the Future Innovation Lab for
Integrated Pest Management



USAID
FROM THE AMERICAN PEOPLE

Tuta absoluta: an invasive alien species

2

Invasive alien species are plants, animals, pathogens and other organisms that are non-native to an ecosystem, and which may cause economic or environmental harm or adversely affect human health.




(UICN 2002)

Why is this pest a threat?

3

The « enemy release » hypothesis




(Keane and Crawley 2002)

Integrated management of *Tuta absoluta*

4

A sound combination of tools...




(adapted from Eilenberg et al. 2001)

Integrated management of *Tuta absoluta*

5

...targeting each stage of the pest.



Monitoring

6

- ✓ Pheromone traps - 2-4 traps/ha and weekly observation
- ✓ Egg/larvae or damage sampling



Spray only when necessary, with as selective as possible insecticides

Cultural control

7

- ✓ Destroy previous crop residue.
- ✓ Remove host weeds near the field or greenhouse.
- ✓ Check the seedlings before transplanting to ensure they are free of eggs and larvae.
- ✓ Remove and destroy any infected leaves, shoots and fruit immediately.



Physical control

8

- ✓ Grow tomato seedlings inside a netted nursery.
- ✓ Double check the net to make sure the greenhouse does not have holes or gaps.
- ✓ Install a secure door in the greenhouse to prevent moths from entering.



Host plant resistance

9

✓ Breeding resistant host plants

Constitutive defenses

Induced defenses

(Biondi et al. 2018)



Biotechnical control

10

- ✓ Mass trapping (20-25 traps/ha greenhouse, 40-50 open field)
- ✓ Mating disruption (30-60 g of pheromone/ha)



(Capparos et al. 2013, Vacas et al. 2011, Cocco et al. 2013)


Biological control

11

The use of living organisms to keep pests below damaging levels .



(Biondi et al. 2018)



Generalist predators

12

Nesidiocoris tenuis and *Macrolophus pygmaeus*
(Hemiptera, Miridae)



©Koppert Biological System



Egg parasitoids

13

Trichogramma achaeae (Hymeno., Trichogrammatidae)



Larval parasitoids

14

Stenomesus japonicus and *Necremnus tuta*
(Hymenoptera, Eulophidae)



Entomopathogens

15

Bacillus thuringiensis var. *kurstaki* or *aizawai*




Fig. 1. Mechanism of toxicity of Bt

Biological control

16

Augmentative

T. achaeae
(inundative)


M. Pygmaeus
N. tenuis
(inoculative)

Classical

.....

Conservation

Banker plants
Trophic resources
Refuges



Towards a system-wide management

17

**ANNUAL
REVIEWS**

Annual Review of Entomology

Ecology, Worldwide Spread, and Management of the Invasive South American Tomato Pinworm, *Tuta absoluta*: Past, Present, and Future

Antonio Biondi,^{1,*} Raul Narciso C. Guedes,²
Fang-Hao Wan,³ and Nicolas Desneux^{4,*}

¹Department of Agriculture, Food and Environment, University of Catania, 95123, Italy; email: antonio.biondi@unict.it

²Departamento de Entomologia, Universidade Federal de Viçosa, Minas Gerais 36570-900, Brazil; email: guedes@ufv.br

³Key Laboratory for Biology of Plant Disease and Insect Pests, Institute of Plant Protection, Chinese Academy of Agricultural Sciences, Beijing 100193, China; email: wanfanghao@caas.cn

⁴INRA (French National Institute for Agricultural Research), University of Côte d'Azur, CNRS, UMR 1355-7254, 06903 Sophia Antipolis, France; email: nicolas.desneux@inra.fr

